arXiv:1411.6996v3 [math.AG] 14 Mar 2015

BOUNDED NEGATIVITY, MIYAOKA-SAKAI INEQUALITY AND ELLIPTIC CURVE CONFIGURATIONS

XAVIER ROULLEAU

Abstract. Similarly to the linear Harbourne constant recently introduced in [1], we study the elliptic H-constants of $\mathbb{P}^2$ and Abelian surfaces. We exhibit configurations of smooth plane cubic curves whose Harbourne index is arbitrarily close to -4. As a Corollary, we obtain that the global H-constant of any surface X is less or equal to -4. Related to these problems, we moreover give a new inequality for the number and multiplicities of singularities of elliptic curves arrangements on Abelian surfaces, inequality which has a close similarity to the one of Hirzebruch for arrangements of lines in the plane.

Introduction

The Harbourne constant (for short H-constant) of a surface and some related variants of it have been recently introduced in [1], bringing new problems and open questions on curves and their singularities on surfaces. As explained there, these constants measure the local negativity of curves on surfaces, in analogy with the local positivity measured by Seshadri constants. These constants come from the context of the Bounded Negativity Conjecture (BNC):

Conjecture. *Let $X_{/\mathbb{C}}$ be a smooth projective surface. There exists an integer $b(X)$ such that for every (reduced) curve C on X, we have $C^2 \geq -b(X)$.*

This ancient and now intensively studied conjecture ([1], [2], [3], [4], [11], [13]) is trivially true for the plane, but we do not know the behavior of the problem if one takes blow-ups of it. The H-constant was introduced to approach that question. For a blow-up $X' \to X$ of a surface X at a set $\mathcal{P}$ of $s > 0$ distinct points and $C \hookrightarrow X$ a curve, we denote by $\bar{C}$ the strict transform of C in X'. The H-constant of X is defined by

$$H_X := \inf_{C,\mathcal{P}} H(C, \mathcal{P})$$

where

$$H(C, \mathcal{P}) = \frac{(\bar{C})^2}{s}$$

for $C \hookrightarrow X$ varying over all reduced curves on X, and the strict transform of C varying over the blows-ups at sets $\mathcal{P}$ of s distinct points on X ($s > 0$ any). If finite, the H-constant has the interesting property that whenever BNC holds for X, then it holds for any blow-ups of it.

Variations of the H-constants are proposed in [1], among them the linear H-constant H_L for the plane. Here $X = \mathbb{P}^2$ and the infimum $H_{L,\mathbb{P}^2} = \inf_{C,\mathcal{P}} H(C, \mathcal{P})$ is taken over every union C of lines in $\mathbb{P}^2$. Using Hirzebruch bounds on the singularities of lines configurations, the authors proves that $H_{L,\mathbb{P}^2} \geq -4$. They moreover found an example of union of lines C such that its Harbourne index

$$H(C) = \min_{\mathcal{P}} H(C, \mathcal{P})$$

is very negative: $H(C) = -225/67$, therefore one knows that $-225/67 \geq H_{L,\mathbb{P}^2} \geq -4$, and $-225/67 \geq H_{L,\mathbb{P}^2} \geq H_{\mathbb{P}^2}$.

In this paper, we similarly propose to study the elliptic H-constant $H_{El,X}$ defined by

$$H_{El,X} = \inf_{C,\mathcal{P}} H(C,\mathcal{P}) = \inf_{C} H(C)$$

for unions C of smooth elliptic curves on the surface X. Let C be such a configuration of elliptic curves on an Abelian surface A. Like for an arrangement of lines in the plane, the curve C has only ordinary singularities. For $k \geq 2$, we denote by t_k be the number of k-points on C i.e. the points with multiplicity k. Let be $f_0 = \sum t_k$ and $f_1 = \sum k t_k$.

Theorem 1. *For any set $\mathcal{P}$ of points, we have*

$$H(C,\mathcal{P}) \geq H(C) = -\frac{f_1}{f_0} \geq \frac{t_2 + \frac{1}{4}t_3}{f_0} - 4 \geq -4 \tag{0.1}$$

The elliptic H-constant satisfies $H_{El,A} \geq -4$. The H-index of C equals -4 if and only if the only singularities on C are 4-points.
The H-constant and elliptic H-constant are isogeny invariants : if A and B are isogenous, then $H_A = H_B$ and $H_{El,A} = H_{El,B}$.

Let A be either the surface $(\mathbb{C}/\mathbb{Z}[\alpha])^2$, where $\alpha^2 + \alpha + 1 = 0$ or $(\mathbb{C}/\mathbb{Z}[i])^2$, $i^2 = -1$. By the constructions of Hirzebruch [7] and Holzapfel [9, Example 5.4], there exist configurations C of elliptic curves on A such that their H-index satisfy $H(C) = -4$. We thus see that the bound $H_{El,A} \geq -4$ is optimal. Actually, by Kobayashi's results [10], the equality $H(C) = -4$ is attained if and only if the complement of C in A is birational to an open ball quotient surface. Inequality

$$H(C) = -\frac{f_1}{f_0} \geq \frac{t_2 + \frac{1}{4}t_3}{f_0} - 4$$

in Theorem 1 is a consequence of a more general result (Theorem 9) on the H-constant of curve configurations of any genus proved by using $(\mathbb{Z}/n\mathbb{Z})^d$-covers of Abelian surfaces. We obtain in particular the following result:

Theorem 2. *For a configuration of elliptic curves on an Abelian surface, one has*

$$t_2 + \frac{3}{4}t_3 \geq \sum_{k\geq 5}(2k-9)t_k.$$

This is the very analog of the inequality $t_2 + \frac{3}{4}t_3 \geq d + \sum_{k\geq 5}(2k-9)t_k$ due to Hirzebruch for a configuration of $d \geq 6$ lines in $\mathbb{P}^2$ such that $t_d = t_{d-1} = t_{d-2} = 0$ (see [8, eq. (9)]).

We also obtain the following results for the elliptic and global H-constants of the plane:

Proposition 3. *There exist configurations C_n of smooth cubic curves in $\mathbb{P}^2$ such that for $Sing(C_n)$ the set of singular points of C_n, we have*

$$\lim_n H(C_n, Sing(C_n)) = -4,$$

and therefore $-4 \geq H_{El,\mathbb{P}^2} \geq H_{\mathbb{P}^2}$.

The configurations C_n appear in [15]. From Proposition 3 and functorial properties of the H-constants, we get the following Corollary on the Harbourne constant of any surface:

Corollary 4. *Let X be a smooth surface. Then $-4 \geq H_{\mathbb{P}^2} \geq H_X$.*

The paper is organized as follows. In the first section, we prove Proposition 3 and Corollary 4 concerning the elliptic H-constant and the Harbourne constant of the plane. In the second section we prove Theorem 2, which together with Theorem 1 proved in the third section are the main results. In the last section we discuss some questions and problems raised by the definitions of the H-index of a curve and H-constant of a surface.

Acknowledgements. The author thanks Piotr Pokora for its relevant remarks on a previous version of this paper.

1. ELLIPTIC CURVE CONFIGURATIONS ON THE PLANE

Let $p : Z \to \mathbb{P}^2$ be the blow-up of $\mathbb{P}^2$ at the 12 singular points $\mathcal{P}_{HE} = \{p_1, \dots, p_{12}\}$ of the dual Hesse configuration. In [15], is described some elliptic curve configurations $\mathcal{H}_n$ ($n \in 3\mathbb{N}$, $n > 0$) on Z with the following properties:
i) $\mathcal{H}_n$ is the union of $\frac{4}{3}(n^2-3)$ elliptic curves which are fibers of some elliptic fibrations of Z,
ii) Each elliptic curve of $\mathcal{H}_n$ has 9 3-points and n^2-9 4-points,
iii) The singularities of $\mathcal{H}_n$ are $\frac{1}{3}(n^2-3)(n^2-9)$ 4-points and $4(n^2-3)$ 3-points.

Let C_n be the image in $\mathbb{P}^2$ of $\mathcal{H}_n$ by the blow-down map p. Then :
i) C_n is the union of $\frac{4}{3}(n^2-3)$ smooth degree 3 curves.
ii) Each elliptic curve contains n^2-9 4-points and goes through 9 of the 12 singular points of the dual Hesse configuration.
iii) The singularities of C_n are $\frac{1}{3}(n^2-3)(n^2-9)$ 4-points and the 12 points in $\mathcal{P}_{HE}$ have multiplicity (n^2-3) (on the blow up at $p_i \in \mathcal{P}_{HE}$, the strict transform of the curve C_n has $\frac{1}{3}(n^2-3)$ 3-points on the exceptional divisor).

Proposition 5. *The configurations C_n of smooth cubic curves on $\mathbb{P}^2$ satisfy*

$$\lim_n H(C_n, \mathcal{P}_n) = -4,$$

where $\mathcal{P}_n$ is the set of singular points of C_n.

Proof. Let be $X_n \to \mathbb{P}^2$ be the blow-up of $\mathbb{P}^2$ at $\mathcal{P}_n$, the $s_n = 12 + \frac{1}{3}(n^2-3)(n^2-9)$ singularities of C_n. Let $\bar{C}_n$ be the strict transform of C_n. The blow down map $X_n \to \mathbb{P}^2$ factorizes through Z. Then

$$\bar{C}_n^2 = \sum \bar{C}_i^2 + 2\sum_{i<j} \bar{C}_i \bar{C}_j$$

where $\bar{C}_i$ is the strict transform of a curve C_i in $\mathcal{H}_n$. Since $\bar{C}_i^2 = -(n^2-9)$ and since from the configuration one has:

$$\sum_{i<j} \bar{C}_i \bar{C}_j = 4(n^2-3),$$

we get

$$\bar{C}_n^2 = -(n^2-9) \times \frac{4}{3}(n^2-3) + 8(n^2-3) = -\frac{4}{3}n^4 + O(n^3).$$

As $s_n = \frac{1}{3}n^4 + O(n^3)$, we obtain the result. □

Remark 6. A) By example, for $n = 7$, we obtain $H(C_n, \mathcal{P}_n) = -\frac{20148}{5257} \simeq -3,83$.
B) The above configurations C_n are strongly linked to some compactifications X_n of some open ball quotient surfaces ([7]), for which $\lim \frac{c_1^2}{c_2}(X_n) = 3$, ie one is close to the upper bound in the Miyaoka-Yau inequality.

Let $f : X \to Y$ be a dominant morphism between two smooth surfaces. Let C be a reduced curve on Y. Suppose that C do not contain components in the branch locus B and let $\mathcal{P}$ be a set of s point in Y disjoint with B (so that f^*C and $f^*\mathcal{P}$ are reduced of pure dimension 1 and 0 respectively).

Lemma 7. *Then $H_X(f^*C, f^*\mathcal{P}) = H_Y(C, \mathcal{P})$.*

Proof. Let d be the degree of f. Let p be a point of $\mathcal{P}$. Then the d points above p have the same multiplicity m_p inside $C' = f^*C$ than p inside C and we have:

$$H_X(f^*C, f^*\mathcal{P}) = \frac{\bar{C'}^2}{ds} = \frac{dC^2 - d\sum m_p^2}{ds} = \frac{\bar{C}^2}{s} = H_Y(C, \mathcal{P}).$$

□

Proposition 5 implies :

Corollary 8. *Let X be a smooth surface. Then $H_X \leq H_{\mathbb{P}^2} \leq -4$.*

Proof. Let $f : X \to \mathbb{P}^2$ be a generic projection of X onto the plane. For any curve C in $\mathbb{P}^2$ and set of point $\mathcal{P} \subset \mathbb{P}^2$, we can choose an automorphism g such that g^*C do not contain any components of the branch divisor B and $g^*\mathcal{P}$ is disjoint with B. We then apply Lemma 7 and results of Proposition 5 to conclude that $H_X \leq H_{\mathbb{P}^2} \leq -4$. □

2. Arrangement of curves on Abelian surfaces and $(\mathbb{Z}/n\mathbb{Z})^d$-covers.

Let A be an abelian surface and let $C = \sum_{i=1}^d C_i$ be a reduced divisor with only ordinary singularities, a union of $d \geq 2$ smooth divisors C_i (by example C_i may be the union of genus 1 fibers of a fibration of A onto an elliptic curve). Let g be the geometric genus of C ie

$$g - 1 = \sum g_j - 1,$$

where the g_j's are the geometric genus of the irreducible components of C. Let us denote by t_k the number of k-points on C.

Theorem 9. *We have*

$$10g - 10 + t_2 + \frac{3}{4}t_3 \geq \sum_{k\geq 5}(2k-9)t_k,$$

and

$$H(C, Sing(C)) = \frac{2g-2-f_1}{f_0} \geq \frac{2t_2 + \frac{9}{8}t_3 + \frac{1}{2}t_4 + 8 - 8g}{f_0} - \frac{9}{2}. \tag{2.1}$$

Suppose that C is a configuration of elliptic curves. We have

$$H(C) = -\frac{f_1}{f_0} \geq \frac{t_2 + \frac{1}{4}t_3}{f_0} - 4 \geq -4 \tag{2.2}$$

where $H(C)$ is the H-index: $H(C) = \min_{\mathcal{P}} H(C, \mathcal{P})$. Moreover $H(C) = -4$ if and only if the only singularities on C are 4-points.

The remaining of this section is the proof of Theorem 9. Let us recall a Theorem of Namba on branched covers. Let M be a manifold, $D_1, \ldots, D_s$ be irreducible reduced divisors on M and $n_1, \ldots, n_s$ be positive integers. We denote by D the divisor $D = \sum n_i D_i$. Let $Div(M, D)$ be the sub-group of the $\mathbb{Q}$-divisors generated by the entire divisors and

$$\frac{1}{n_1}D_1, \ldots, \frac{1}{n_s}D_s.$$

Let $\sim$ be the linear equivalence in $Div(M, D)$, where $G \sim G'$ if and only if $G - G'$ is an entire principal divisor. Let $Div(M, D)/\sim$ the quotient and let $Div^0(M, D)/\sim$ be the kernel of the Chern class map

$$\begin{array}{ccc} Div(M,D)/\sim & \to & H^{1,1}(M, \mathbb{R}) \\ G & \to & c_1(G) \end{array}.$$

Theorem 10. *(Namba,* [14, Thm. 2.3.20]*). There exists a finite Abelian cover which branches at D with index n_i over D_i for all $i = 1, \ldots, s$ if and only if for every $j = 1, \ldots, s$ there exists an element of finite order $v_j = \sum \frac{a_{ij}}{n_i} D_i + E_j$ of $Div^0(M, D)/\sim$ (E_j an entire divisor, $a_{ij} \in \mathbb{Z}$) such that a_{jj} is coprime to n_j.*
Then the group in $Div^0(M, D)/\sim$ generated by the v_j is isomorphic to the Galois group of one of such Abelian cover.

We find the inequalities among the t_k's in Theorem 9 using $(\mathbb{Z}/n\mathbb{Z})^d$-covers of A ramified above curves (related to the) C_i. These inequalities involve quantities that are "linear" under isogenies, by which we mean that if $\phi : B \to A$ is an isogeny of degree m, then the number of k-points on ϕ^*C (a reduced curve), the intersections between the ϕ^*C_i's and ϕ^*C, the geometric genus minus 1 of ϕ^*C etc... are the ones of C, C_i etc... multiplied by m. By that

property, inequalities involving linear terms in the t_k's and C_i^2 proved on abelian surface B are then inequalities for A.

Recall that if $\phi = [m] : A \to A$ is the multiplication by $m \in \mathbb{N}$ map, then $\phi^* D \sim \frac{m(m+1)}{2} D + \frac{m(m-1)}{2}[-1]^* D$ for any divisor D. By taking $m = 2n$, one can therefore suppose that the divisors C_i are n-divisible ie there exists divisors L_i such that $C_i \sim nL_i$. The divisor $v_i = \frac{1}{n}C_i - L_i$ is in $Div^0(A, nC)/\sim$, has order n and the multiplicity of an irreducible component C_i' in $\frac{1}{n}C_i$ is $\frac{1}{n}$. The group generated by divisors $\frac{1}{n}C_i - L_i$ is isomorphic to $(\mathbb{Z}/n\mathbb{Z})^d$ and there exists a $(\mathbb{Z}/n\mathbb{Z})^d$-cover of A branched with index n over C. For the computation of the Chern numbers of the resolution X_n of that cover we refer to the local analysis of the $(\mathbb{Z}/n\mathbb{Z})^d$-branched covers of the plane constructed by Hirzebruch in [7] (see also the geometric approach of [5]).

The following quantities f_0, f_1, f_2 are linear under isogenies:

$$f_0 = \sum_{k\geq 2} t_k,\ f_1 = \sum_{k\geq 2} k t_k,\ f_2 = \sum_{k\geq 2} k^2 t_k.$$

Let $\pi : Z \to A$ be the blow-up at the $f_0 - t_2 = \sum_{k\geq 3} t_k$ singularities of C of multiplicities $k \geq 3$ and let $\bar{C} = \sum \bar{C}_i$ be the strict transform of C in Z. For a singularity p of C of multiplicity $k_p \geq 3$, we denote by $E_p \hookrightarrow Z$ the exceptional curve over p. There exists a degree n^d map

$$f : X_n \to Z$$

branched with index n above the curve $\bar{C}$. Above E_p lies n^{d-r} $(r = k_p)$ copies in X_n of a smooth curve F_p which is a $(\mathbb{Z}/n\mathbb{Z})^{r-1}$-cover of E_p ramified with index n at r points, thus

$$e(F_p) = n^{r-1}(2-r) + rn^{r-2} = n^{r-2}(2n + r(1-n)).$$

Since the Galois group permutes these curves, we have $(F_p)^2 = -n^{r-2}$. If a singularity p of C is a node, then X_n is smooth over p and the fiber of f at p has only n^{d-2} points.

We have

$e(C) = 2 - 2g + f_0 - f_1$, $e(C \setminus Sing(C)) = 2 - 2g - f_1$, $e(A \setminus C) = -e(C) = 2g - 2 + f_1 - f_0$, and $C^2 = \sum C_i^2 + f_2 - f_1 = 2g - 2 + f_2 - f_1$. Therefore we obtain

$$e(X_n \setminus f^{-1}E_p) = n^d e(A \setminus C) + n^{d-1} e(C \setminus Sing(C)) + n^{d-2} t_2$$

and

$$e(X_n \setminus f^{-1}E_p)/n^{d-2} = n^2(2g - 2 + f_1 - f_0) + n(2 - 2g - f_1) + t_2.$$

Since above each exceptional divisor E_p in Z, we have n^{d-k} curves with Euler number $e(F_p)$, we get

$$e(X_n) = e(X_n \setminus f^{-1}E_p) + \sum_{k\geq 3} n^{d-2} t_k (2n + k(1-n))$$

thus

$$e(X_n)/n^{d-2} = (2g - 2 + f_1 - f_0)n^2 + 2(1 - g + f_0 - f_1)n + f_1 - t_2.$$

Let us now compute the canonical divisor : K_{X_n} is numerically equivalent to the pullback of

$$K = \sum E_p + \frac{n-1}{n}(\sum E_p + \pi^* C - \sum k_p E_p) = \sum_p \frac{2n - 1 + k_p(1-n)}{n} E_p + \frac{n-1}{n}\pi^* C,$$

thus

$$K^2 = \sum_{k\geq 3} -\frac{(2n - 1 + k(1-n))^2}{n^2} t_k + (\frac{n-1}{n})^2 C^2.$$

We obtain

$$K_{X_n}^2/n^{d-2} = (2g - 2 + 3f_1 - 4f_0)n^2 + 4(f_0 - f_1 - g + 1)n - f_0 + f_1 + t_2 + 2g - 2.$$

Since it covers an Abelian surface the Kodaira dimension of X_n is non negative. Then we get by using the Miyaoka-Yau inequality

$$(3c_2 - K^2_{X_n})/n^{d-2} = (f_0 + 4g - 4)n^2 + 2(f_0 - f_1 - g + 1)n + 2f_1 + f_0 - 4t_2 - 2g + 2 \geq 0. \quad (2.3)$$

As in [8], we will use refinements of the Miyaoka-Yau inequality for surfaces that contain smooth rational curves and elliptic curves, ie for $n = 2$ or 3. Let Y be a surface of non negative Kodaira dimension. Suppose that there exists on Y smooth disjoint elliptic curves D_j and m disjoint (-2)-curves, disjoint also from the curves D_j, then:

Theorem 11. *(Miyaoka* [12, Cor. 1.3]*). We have*

$$3c_2(Y) - K^2_Y \geq \frac{9}{2}m - \sum (D_j)^2.$$

For $n = 3$, we get from equation 2.3:

$$(3c_2 - K^2_{X_3})/3^{d-2} = 4(7g - 7 + 4f_0 - f_1 - t_2) \geq 0$$

Taking into account the fact that over the 3-points the surface contains $3^{d-3}t_3$ elliptic curves of self-intersection -3, we can refine that inequality and obtain

$$-\frac{f_1}{f_0} \geq \frac{t_2 + \frac{1}{4}t_3 - 7g + 7}{f_0} - 4.$$

For $g = 1$, ie C is a configuration of elliptic curves, let $Sing(C)$ be the singularity set of C, then:

$$H(C, Sing(C)) = -\frac{f_1}{f_0} \geq \frac{t_2 + \frac{1}{4}t_3}{f_0} - 4.$$

For $\mathcal{P}$ a (non void) set of points in A, one can use the same demonstration as in [1, thm 3.3] for the linear H-constant of $\mathbb{P}^2$ to conclude that

$$H(C, \mathcal{P}) \geq H(C) = -\frac{f_1}{f_0} \geq \frac{t_2 + \frac{1}{4}t_3}{f_0} - 4.$$

Suppose that the bound -4 is attained, then $4f_0 = f_1$, $t_2 = t_3 = 0$ and from equality $\sum_{k\geq 5}(k-4)t_k = 2t_2 + t_3$, we see that $t_k = 0\ \forall k \geq 5$. The only possibility is $t_4 \neq 0$, which indeed exists (see below).

For $n = 2$, the surface X_2 contains $2^{d-3}t_3$ disjoint (-2)-curves and it contains $t_4 2^{d-4}$ elliptic curves of self-intersection -4, therefore

$$(3c_2 - K^2_{X_2})/2^{d-2} \geq \frac{9}{4}t_3 + t_4$$

and

$$10g - 10 + 9f_0 \geq 2f_1 + 4t_2 + \frac{9}{4}t_3 + t_4.$$

We obtain the following inequality:

$$10g - 10 + t_2 + \frac{3}{4}t_3 \geq \sum_{k\geq 5}(2k-9)t_k.$$

Using $H(C, Sing(C)) = \frac{C^2 - f_2}{f_0} = \frac{\sum C_i^2 - f_1}{f_0}$ and $2g - 2 = \sum C_i^2$, we obtain

$$H(C, Sing(C)) \geq \frac{2t_2 + \frac{9}{8}t_3 + \frac{1}{2}t_4 + 8 - 8g}{f_0} - \frac{9}{2}.$$

3. The Elliptic H-constants for Abelian surfaces

Let X be a smooth projective surface and C be a configuration of smooth disjoint elliptic curves on X. Let us recall the following result:

Theorem 12. *(Kobayashi) There exists a torsion free lattice $\Gamma \subset PU(2,1)$ such that (X, C) is a smooth compactification of the ball quotient surface $\mathbb{B}_2/\Gamma$ if and only if $(K_X + C)^2 = 3e(X \setminus C)$.*

Let C be an elliptic curve configuration on a an Abelian surface A. Let $X \to A$ be the blow-up at the singular points of C and let $\bar{C}$ be the strict transform of C in X. We obtain:

Corollary 13. *The elliptic curve configuration C on A has H-index $H(C) = -4$ if and only if $(X, \bar{C})$ is a smooth compactification of a ball quotient surface.*
In that case, the curve C has only 4-points singularities and there exists a covering $X_3 \to A$ branched with order 3 over $[3]^*C$ such that X_3 is a smooth compact ball quotient surface: $c_1^2(X_3) = 3c_2(X_3)$.

Remark 14. Hirzebruch constructed such compact ball quotient surface in [7] for the case of the Eisenstein Abelian surface (see below).

Proof. One computes that $(K_X + \bar{C})^2 = f_1 - f_0$ and $e(X \setminus \bar{C}) = f_0$. Therefore, one has equality $(K_X + \bar{C})^2 = 3e(X \setminus \bar{C})$ if and only if $4f_0 = f_1$, which is equivalent by Theorem 9 to $H(C) = -4$ and to the condition that C has only 4-points.

By equation 2.3, the value of $(3c_2(X_n) - K_{X_n}^2)/n^{d-2}$ is $f_0(n-3)^2$, thus for $n = 3$, we obtain a compact ball quotient surface. □

Let be $j = \frac{-1+i\sqrt{3}}{2}$ with $i^2 = -1$, we have:

Proposition 15. *The elliptic H-constants of $(\mathbb{C}/\mathbb{Z}[i])^2$ and $(\mathbb{C}/\mathbb{Z}[j])^2$ are equal to -4.*

Proof. Hirzebruch [7] and Holzapfel [9] found elliptic curves arrangements on $(\mathbb{C}/\mathbb{Z}[i])^2$ and $(\mathbb{C}/\mathbb{Z}[j])^2$ respectively that satisfies Kobayashi's criteria on suitable blow-up. In the first example, one has 4 elliptic curves with $t_4 = 1$ and $t_k = 0$, $k \neq 4$, in the second one has 6 elliptic curves with $t_4 = 3$ and $t_k = 0$, $k \neq 4$. □

Remark 16. Hirzebruch and Holzapfel examples are elliptic curves configurations on Abelian surfaces with only 4-points singularities. That situation must be compared with the plane where there do not exist configurations of $d > 6$ lines with $t_2 = t_3 = 0$ and $t_d = t_{d-1} = t_{d-2}$ (by inequality 4.3 below).

Let E, E' be two elliptic curves and let A be an Abelian surface.

Proposition 17. *The H-constant and elliptic H-constant of A are invariants of the isogeny class of A. Suppose A is isogenous to $E \times E'$. If E and E' are not isogenous, then $H_{El,A} = -2$. If E and E' are isogenous, then $H_{El,A} \leq -3$.*

Proof. Let $\phi : A \to B$ be an isogeny between two Abelian surfaces ; it is an étale map. Let C be a (reduced) curve on B, and let $\mathcal{P}$ be a set of points on B. By Lemma 7, $H_A(\phi^*\mathcal{P}, \phi^*C) = H_B(C, \mathcal{P})$, thus

$$\inf_{C,\mathcal{P}} H_A(C, \mathcal{P}) \leq \inf_{C,\mathcal{P}} H_B(C, \mathcal{P}).$$

Since there exists an isogeny $\psi : B \to A$ too, we have the reverse inequality. That holds also for the elliptic H-constant, since the pull-back by an isogeny of a genus one curve is a union of genus one curves.

Let A be isogenous to $E \times E'$. Suppose that E and E' are not isogenous. Then a configuration C of elliptic curves on $E \times E'$ is as follows:

$$C = \sum_{k=1}^{m} F_k + \sum_{k=1}^{n} F'_k$$

where the F_k (resp. F'_k) are fibers of the fibration of $E \times E'$ onto E (resp. E'). Then one sees that $\bar{C}^2/s$ is minimal and equals -2 when the blow-up is taken over the mn intersection points of F_i and F'_j.

Suppose that E and E' are isogenous. Since the elliptic H-constant is isogeny invariant, we can suppose that $E = E'$. Let Δ be the diagonal in $E \times E$ and let be $F = \{y = 0\}$, $F' = \{x = 0\}$, where x, y are the coordinates. Then $C = \Delta + F + F'$ has one 3-point in 0, and in the blow-up at 0, we get $\bar{C}^2 = -3$. □

4. Remarks on the Harbourne index of curves

4.1. **Irreducible curves with low H-index.** In view of BNC Conjecture , it would be interesting to know irreducible curves C with low Harbourne index

$$H(C) = \min_{\mathcal{P}} H(C, \mathcal{P}),$$

then by taking the appropriate blow-up one could get very negative curves and maybe obtain counter examples to that Conjecture. By example, if one consider the rational curves of degree d in $\mathbb{P}^2$ with the maximal number of nodes, the H-index $H(C)$ is close to -2 (on that subject, see also Introduction of [1]). But if one is willing to go down, we must impose singularities with higher multiplicities on C, and it forces C to have "negative genus" ie to be union of (at least) two curves. This explain the idea to consider union of curves rather than irreducible curves. Moreover considering reducible curves gives more functorialities to the H-constants, e.g. when one wishes to compare these constants through a dominant map $f : X \to Y$.

On that problem of constructing infinitely many curves with low H-index, the example of totally geodesic curves on a Shimura surface X is particularly interesting. If smooth, such a curves C satisfy $C^2 = -2(g-1)$ (for g the genus) and were considered as possible counter examples for BNC. But one of the main results of [2] is the fact on X, there are at most a finite number of such curves C with $C^2 < 0$. However, when one take some blow-up of X, one do not know the behavior of $\bar{C}^2$ for $\bar{C}$ the strict transform of such a curve C.
The singularities of a totally geodesic curve C on a Shimura or a Ball quotient surface are ordinary, and so are the singularities of a union of such curves. This follows from the fact that C is the image of the trace of lines in the universal cover, and singularities of a union of lines are ordinary. Let X be a Shimura surface and C be a (irreducible) totally geodesic curve on it. Let $\delta \in \mathbb{N}$ be defined by $K_X C + C^2 = 2g - 2 + 2\delta$. We have $2\delta = f_2 - f_1$ (where $f_i = \sum k^i t_k$, for t_k the number of k-points). Since C is a totally geodesic curve, we have $K_X C = 4g - 4 > 3g - 3$ and therefore [4, Cor. 4.2] implies that $C^2 \sim 2\delta$ when the geometric genus g of C goes to ∞. Therefore

$$H(C, Sing(C)) \sim -\frac{f_1}{f_0}$$

when $g \to \infty$ and since always $f_1 \geq 2f_0$, we obtain a second example of irreducible curves on a surface with

$$\liminf_{C} H(C) \leq -2.$$

4.2. **The H-index of line arrangement.** For the case of an arrangement of d lines in $\mathbb{P}^2$ with $t_d = t_{d-1} = t_{d-2} = t_{d-3} = 0$ and $d \geq 6$, Hirzebruch [6, 140] proved the following inequality

$$t_2 + \frac{3}{4}t_3 \geq d + \sum_{k\geq 5}(k-4)t_k. \tag{4.1}$$

Using that results, the authors of [1] obtained the following inequality for the H-index of such a line arrangement C:

$$H(C) \geq B_1 := -4 + \frac{1}{\sum t_k}(2d + t_2 + \frac{1}{4}t_3). \tag{4.2}$$

A better inequality for the singularities of line arrangements is given in [8, eq. (9)], which is

$$t_2 + \frac{3}{4}t_3 \geq d + \sum_{k\geq 5}(2k-9)t_k. \tag{4.3}$$

Proposition 18. *Using inequality 4.3, we obtain:*

$$H(C) \geq B_2 := \frac{1}{\sum t_k}(\frac{3}{2}d + 2t_2 + \frac{9}{8}t_3 + \frac{1}{2}t_4) - \frac{9}{2}. \tag{4.4}$$

and moreover that inequality 4.4 is sharper than 4.2 : $B_2 \geq B_1$.

For the Klein configuration of lines (see [1, § 4.1]) and the Fermat configuration of 18 lines, inequality 4.4 is even an equality.

Proof. Using the combinatorics, we have

$$H(C, Sing(C)) = \frac{d^2 - \sum_{k\geq 2} k^2 t_k}{\sum t_k} = \frac{d - \sum_{k\geq 2} k t_k}{\sum t_k}.$$

By 4.3, we have $t_2 + \frac{3}{4}t_3 \geq d + \sum_{k\geq 4}(2k-9)t_k$, thus

$$-\sum_{k\geq 2} k t_k \geq \frac{1}{2}(d + 4t_2 + \frac{9}{4}t_3 + t_4 - 9\sum t_k) \tag{4.5}$$

and we obtain inequality 4.4. The fact that this new inequality is sharper comes from the fact that inequality 4.3 is better than 4.1. □

References

[1] Bauer T., Di Rocco S., Harbourne B., Huizenga J., Lundman A., Pokora P., Szemberg T., *Bounded negativity and arrangement of lines*, to appear in Int. Math. Res. Notices.

[2] Bauer T., Harbourne B., Knutsen A.L.; Küronya A., Müller-Stach S., Roulleau X., Szemberg T., *Negative curves on algebraic surfaces*, Duke Math. J. 162 (2013), 1877–1894.

[3] Bauer T., Pokora P., Schmitz D., *On the boundedness of the denominators in the Zariski decomposition on surfaces*, arXiv 1411.2431

[4] Ciliberto C., Roulleau X., *On finiteness of curves with high canonical degree on a surface*, arXiv:1406.7478

[5] Gao Y., *A note on finite abelian covers*, Sci. China Math. 54 (2011), no. 7, 1333–1342.

[6] Hirzebruch F., *Arrangement of lines and Algebraic surfaces*, Arithmetic and geometry, Vol. II, 113–140, Progr. Math., 36, Birkhäuser, Boston, Mass., 1983.

[7] Hirzebruch F., *Chern numbers of algebraic surfaces: an example*, Math. Ann. 266 (1984), no. 3, 351–356.

[8] Hirzebruch F., *Singularities of algebraic surfaces and characteristic numbers*, The Lefschetz centennial conference, Part I (Mexico City, 1984), 141–155, Contemp. Math., 58, 1986.

[9] Holzapfel R.-P., *Jacobi Theta Embedding of a Hyperbolic 4 Space with Cusps*, in Geometry, Integrability and Quantization, I. Mladenov and G. Naber (Eds), Corall Press, Sofia 2001, pp.11-63.

[10] Kobayashi R., *Uniformization of complex surfaces*, Kahler metric and moduli spaces, 313–394, Adv. Stud. Pure Math., 18-II, Academic Press, Boston, MA, 1990.

[11] Koziarz V., Maubon J., *On the equidistribution of totally geodesic submanifolds in compact locally symmetric spaces and application to boundedness results for negative curves* , arXiv: 1407.6561

[12] Miyaoka Y., *The maximal number of quotient singularities on surfaces with given numerical invariants*, Math. Ann. 268 (1984), no. 2, 159–171.
[13] Moeller M., Toledo D., *Bounded negativity of self-intersection numbers of Shimura curves on Shimura surfaces* arXiv:1407.5181
[14] Namba, M., *Branched coverings and algebraic functions*, Pitman Research Notes in Mathematics Series, 161. John Wiley Sons, Inc., New York, 1987. viii+201 pp.
[15] Roulleau X., Urzua G., *Chern slopes of simply connected complex surfaces of general type are dense in* $[2, 3]$, ArXiv 1404.3074, to appear in Annals of Math.
[16] Sakai F., *Semistable curves on algebraic surfaces and logarithmic pluricanonical maps*, Math. Ann. 254 (1980), no. 2, 89–120.

Xavier Roulleau,
Laboratoire de Mathématiques et Applications, Université de Poitiers,
Téléport 2 - BP 30179 - 86962 Futuroscope Chasseneuil, France
roulleau@math.univ-poitiers.fr